

# Diabetes and Risk of Non-Hodgkin's Lymphoma

## A meta-analysis of observational studies

JOANNA MITRI, MD<sup>1</sup>  
 JORGE CASTILLO, MD<sup>2</sup>  
 ANASTASSIOS G. PITTAS, MD, MS<sup>3</sup>

**OBJECTIVE** — To examine the epidemiologic association between diabetes and risk of non-Hodgkin's lymphoma (NHL).

**RESEARCH DESIGN AND METHODS** — We searched MEDLINE for observational studies on the association between diabetes and NHL in adults using the keywords “diabetes” and “lymphoma.” Prospective cohort studies that reported relative risks or standardized incidence ratios and case-control studies that reported odds ratios with 95% CIs were included. A random-effects model was used to combine results from the individual studies.

**RESULTS** — A total of 15 manuscripts (reporting data from 5 prospective cohort and 11 case-control studies) met the inclusion criteria. Combining data from all studies, the risk ratio (RR) of developing NHL in patients with diabetes was 1.19 (95% CI 1.04–1.35). Based on prospective studies, patients with diabetes had an RR of developing NHL of 1.41 (1.07–1.88), without heterogeneity among studies ( $I^2 = 34.3\%$ ;  $P > 0.10$ ). Based on case-control studies, patients with diabetes had an RR of 1.12 (95% CI 0.95–1.31) of developing NHL compared with people without diabetes, with some heterogeneity among studies ( $I^2 = 36.28\%$ ;  $P = 0.09$ ).

**CONCLUSIONS** — Diabetes is associated with a moderately increased risk of NHL, which is consistent with other reported associations between diabetes and malignancies. Future studies should focus on elucidating potential pathophysiologic links between diabetes and NHL.

*Diabetes Care* 31:2391–2397, 2008

The incidence of non-Hodgkin's lymphoma (NHL), a heterogeneous group of conditions characterized by uncontrolled growth of monoclonal malignant lymphocytes, has been increasing in Western countries over the last few decades (1). The increase has been more pronounced in whites, men, and the elderly (1). NHL is now the fifth most commonly diagnosed cancer among both men and women and accounts for 4% of all cancer diagnoses (1). Several hypotheses have been proposed to explain this increased incidence, including aging of the population, exposure to pesticides, prolonged survival of patients with immuno-

deficiency states (including HIV), and increased prevalence of autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Sjögren's Syndrome, and multiple sclerosis).

In parallel with the increasing incidence of NHL, the prevalence of diabetes has increased over the last few decades (2). Diabetes has been associated with increased risk of several malignancies, including endometrial, ovarian, breast, gallbladder, liver, colonic, and pancreatic cancer (3–7), but its relationship with NHL is unclear. Individuals with diabetes are believed to have altered immune function and to be in a relatively immuno-

deficient state (8), which may contribute to the pathogenesis of lymphoproliferative disorders, such as NHL (9). Furthermore, there is a growing body of evidence linking insulin and IGF-I to cell proliferation, lack of apoptosis in cells, and cell metastasis (10)—phenomena involved in the development and progression of malignancies.

An association between diabetes and NHL was initially reported 40 years ago (11). Since then, several studies have examined this association in different population- and hospital-based settings with inconsistent results. The objective of this study was to perform a systematic review and meta-analysis of observational studies to estimate the risk of NHL in patients with diabetes.

## RESEARCH DESIGN AND METHODS

### Data search

We conducted a search of MEDLINE (1 January 1980 through 30 April 2008) for observational cohort and case-control studies reporting an association between diabetes and NHL using the keywords “diabetes” and “lymphoma.” We searched for additional studies in personal reference lists and citation sections of recovered articles.

### Study selection

Two authors (J.M. and J.C.) screened abstracts according to the inclusion criteria. An abstract was judged relevant if it reported original data, was published in the English language, and was from epidemiological studies where the outcome variable was NHL and the predictor variable was diabetes (any type). Prospective cohort studies reporting relative risks or standardized incidence ratios with 95% CIs and case-control studies reporting odds ratios (ORs) with 95% CIs were included in the meta-analysis. Manuscripts reporting only data on Hodgkin's disease and multiple myeloma were excluded because the pathophysiology of these conditions is different from that of NHL (12). If a manuscript included data on other

From the <sup>1</sup>Department of Medicine, Roger Williams Hospital, Boston University School of Medicine, Providence, Rhode Island; the <sup>2</sup>Division of Hematology and Oncology, Miriam Hospital, Brown University Alpert Medical School, Providence, Rhode Island; and the <sup>3</sup>Division of Endocrinology, Diabetes and Metabolism, Tufts Medical Center, Tufts University School of Medicine, Boston, Massachusetts.

Corresponding author: Anastassios G. Pittas, apittas@tuftsmedicalcenter.org.

Received 7 June 2008 and accepted 20 August 2008.

DOI: 10.2337/dc08-1034

© 2008 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

cancers besides NHL, we extracted the data on NHL only. Full-text articles, including letters to the editor, were retrieved and reviewed if a decision on inclusion could not be made based solely on the abstract. Any discrepancies were resolved by consensus in-group conference referencing the original article. When there were multiple publications from the same population, we included data from the most recent one. We did not contact the authors of the original manuscripts for additional data.

**Data extraction**

The following data were extracted from each study: author, country of origin, publication year, participant characteristics (age and sex), sample size, inclusion and exclusion criteria, and methods of diagnosis of diabetes and NHL. For cohort studies, we extracted the source of cohort, the follow-up period, the source of the expected incidence of NHL, relative risk or standardized incidence ratio with 95% CI, and variables adjusted for in the statistical analysis. For case-control studies, we extracted the source of case and control subjects (e.g., hospital based and population based), OR with 95% CI, and variables adjusted for in the statistical analysis. Ascertainment of exposure (self-reported versus registry based) and extent of multivariate adjustment were used to evaluate study quality. Data were extracted by one author (J.M.) and independently verified by another (A.G.P.).

**Data synthesis and analysis**

The primary measure was risk ratio (RR) of NHL, calculated using the random effects model (DerSimonian-Laird method), which accounts for heterogeneity among studies (13). Because the absolute risk of NHL is low, the OR in case-control studies mathematically approximates the RR (14); therefore, we report all results as RRs. To assess heterogeneity among the studies, we used the Cochran Q and  $I^2$  statistics; for the Q statistic, a  $P$  value  $<0.10$  was considered statistically significant for heterogeneity (15); for  $I^2$ , a value  $>50\%$  is considered a measure of severe heterogeneity (16). Subgroup analyses were performed for sex and for type of diabetes. We assessed publication bias using the funnel plot, which plots relative risk (or OR) by sample size. When data were not uniformly reported to allow formal statistical analyses, we present the data in a narrative format. Data were analyzed using MIX version 1.7 (17), a




Figure 1—Search results.

comprehensive free software for meta-analysis of causal research data that has been validated against STATA (StataCorp; College Station, TX) and Comprehensive Meta-analysis (Biostat; Englewood, NJ), two of the most utilized meta-analysis software tools.

**RESULTS**

**Search results**

A total of 540 abstracts were identified during the initial search (Fig. 1). After review of the abstracts, 514 were ineligible because they were reviews, editorials, case reports, comments, or preclinical studies. After reviewing the reference list of the remaining 26 studies, 8 more studies were considered. After detailed evaluation of the 34 potential manuscripts, 19 manuscripts were excluded for reasons shown in Fig. 1.

**Characteristics of prospective studies**

We identified five prospective cohort studies of diabetes and risk of NHL (Table 1) (5,6,18–20). Studies were published between 1997 and 2006 and included more than 160,000 participants with dia-

betes; participants were followed for 6 to 25 years, reporting a total of 337 incident cases of NHL. The diagnosis of diabetes was self-reported in two studies (6,18) and was registry based in three studies (5,19,20). All studies included histologically confirmed cases of NHL. Two studies excluded NHL cases that were reported during the first 1 or 2 years of follow-up to minimize the impact of selection bias (6,20). None of the studies mentioned whether participants with HIV infection were excluded.

Three studies reported relative risk (6,18), whereas the remaining three reported standardized incidence ratio (5,19,20). With the exception of two studies (6,18), investigators did not adjust risk for potential confounders other than age and sex. One study was performed in a women-only cohort (18).

**Characteristics of case-control studies**

We identified 10 manuscripts reporting data from 11 case-control studies published between 1988 and 2007 (Table 2) (7,21–29). A total of 382 cases of NHL were reported among 4,844 patients with diabetes, whereas a total of 7,974 cases of

**Table 1—Prospective cohort studies of diabetes and risk of NHL**

Authors, year (country)	Cohort	Age (years)	Sex	Type of diabetes	NHL assessment	Mean follow-up (starting to ending year)	n with diabetes/ person-years/ n NHL cases	Definition of diabetes	Adjustments	Study quality
Cerhan et al., 1997 (U.S.)	Iowa Women's Health Study	55–69	F	Type 1 (onset ≤30 years)	National cancer registry	7 (1986–1992)	NR/603/1	Self-reported	Age adjusted	
Hjalgrim et al., 1997 (Denmark)	Funen County, Denmark	NR	M/F	Insulin-treated type 2 (onset ≥30 years)	National cancer registry	19 (1973–1992)	772/7,802/6	Prescriptions based	No adjustments	
Weidenpass et al., 1997 (Sweden)	Swedish National Inpatient Register	NR	M/F	NR	National cancer registry	24 (1965–1989)	134,098/901,147/237	Hospital discharge diagnosis	No adjustments; no difference according to sex	
Swerdlow et al., 2005 (U.K.)	Diabetes UK cohort	NR	M/F	Insulin-treated type 1 (onset <30 years)	National cancer registry	18 (1972–2003)	23,834/NR/14	Registry based	No adjustments	
		NR	M/F	Insulin-treated type 2 (onset 30–49 years)	National cancer registry	18 (1972–2003)	5,066/NR/12	Registry based	No adjustments	
Khan et al., 2006 (Japan)	Japan Nationwide Cohort	40–80	M	NR	Self-reported (confirmed by cancer registry in about half of cohort)	9 (1988–1997)	1,753/NR/28	Self-reported	Adjusted for age, BMI, smoking, and alcohol use	
		40–80	F	NR	Self-reported (confirmed by cancer registry in about half of cohort)	9 (1988–1997)	1,541/NR/19	Self-reported	Adjusted for age, BMI, smoking, and alcohol use	

NR, not reported.

NHL were reported among 86,660 control subjects without diabetes. The diagnosis of diabetes was self-reported in all studies. The diagnosis of NHL was pathologically confirmed in all studies with the exception of one study that confirmed the diagnosis of NHL through a review of medical records (27). Control subjects originated from a population-based (7,21,22,26–29) or hospital-based setting (23–25). All studies adjusted their result for age. Eight studies adjusted for sex (21,23–29), two for BMI (7,24), three for ethnicity (7,21,28), two for level of education (7,25), and one for country of origin (29). Four studies excluded individuals with HIV infection (21,24,27,29). Two studies included only men (7,22).

**Analyses**

After combining data from all studies (prospective and case-control), the RR of NHL in patients with diabetes was 1.19 (95% CI 1.04–1.35) (Fig. 2). Visual examination of the funnel plot revealed minimal asymmetry. In prospective studies, the RR for NHL among patients with diabetes was 1.41 (95% CI 1.07–1.88), without heterogeneity among studies ( $I^2 = 34.3\%$ ;  $P > 0.10$ ). Among case-control studies, the increased risk of NHL among patients with diabetes was not statistically significant (1.12 [0.95–1.31]) but had a mild degree of heterogeneity ( $I^2 = 36.28\%$ ,  $P = 0.09$ ).

**Diabetes and NHL by sex**

We then combined data from the three prospective (6,18,20) and four case-control (7,22,25,28) studies that reported risk of NHL according to sex. Female patients with diabetes had a statistically significant increased risk of developing NHL (RR 1.38 [95% CI 1.06–1.80]). In men with diabetes, there was no increased risk of NHL (0.98 [0.79–1.22]). No heterogeneity was detected in these subgroup analyses.


**Type of diabetes and NHL**

Along with one study that identified type of diabetes by national registry (5), several studies used the age of onset of diabetes (18–21,24,29) as a proxy for a presumptive diagnosis of type 1 or type 2 diabetes (Tables 1 and 2). Combining databases from patients with type 1 diabetes (age of onset younger than 30 years), the pooled RR was 1.27 (95% CI 0.82–1.99), without evidence of heterogeneity ( $I^2 = 0.0\%$ ;  $P > 0.10$ ). In patients with type 2 diabetes (age of onset older than 30

Table 2—Case-control studies of prevalence of diabetes among patients with NHL

Authors, year (country)	Sex	Age (years)	Cases (ascertainment period), n	Control subjects, n	Exposure, type of diabetes	Study Quality	
						Exposure, ascertainment method	Adjustments
Bernstein et al., 1988 (U.K.)	M/F	19–75	Incident NHL (1979–1982), 619	Population based, 619	Type NR	Self-reported	Age, sex, race, residence
Cartwright et al., 1988 (U.K.)	M/F	NR	NHL (1979–1984), 458	Hospital based, 742	Type NR	Self-reported	Age, sex, residence
Zham et al., 1995 (U.S.) Kansas cohort	M	NR	NHL (1976–1982), 169	Population based, 936	Type NR	Self-reported	Age
Zham et al., 1995 (U.S.) Iowa/Minnesota cohort	M	NR	NHL (1980–1983), 577	Population based, 1,146	Type NR	Self-reported	Age
Vineis et al., 2000 (Italy)	M/F	20–74	NHL (1990–1993), 1,388	Population based, 1,718	Type NR	Self-reported	Age, sex
Cerhan et al., 2005 (U.S.)	M/F	20–74	Incident NHL (1998–2000), 703	Population based, 538	Type 1 (onset ≤ 30 years)	Self-reported	Age, sex, ethnicity, study site. HIV+ excluded
Fortuny et al., 2005 (Spain)	M/F	17–96	Incident NHL (1998–2000), 753	Population based, 587	Type 2 (onset > 30 years)	Self-reported	Age, sex, ethnicity, study site. HIV+ excluded
Rousseau et al., 2005 (Canada)	M	35–70	Incident NHL (1979–1985), 195	Population based, 509	Type NR	Self-reported	Age, BMI, ethnicity, income, education, farming
Smedby et al., 2006 (Denmark and Sweden)	M/F	17–96	Incident NHL (B-cell) (1998–2002), n = 247	Hospital based, 595	Type 2 (onset > 30 years)	Self-reported	Age, sex, study center. HIV+, post-transplant excluded
	M/F	17–96	Incident NHL (T-cell) (1998–2002), 45	Hospital based, 595	Type 2 (onset > 30 years)	Self-reported	Age, sex, study center. HIV+, post-transplant excluded
Lin et al., 2007 (Taiwan)	M/F	>30	Incident NHL (2000–2004), 242	Population based, 71,379	Type NR	Self-reported	Age, sex, smoking, alcohol, hypertension. HIV+ excluded
Scotti et al., 2007 (Italy)	M/F	17–85	NHL (2000–2004), 671	Hospital based, 1,799	Type NR	Self-reported	Age, sex, study site, residence, education

NR, not reported.



**Figure 2**—Risk estimates of developing NHL for patients with diabetes compared with control subjects (patients without diabetes).

years), the pooled RR was 1.32 (0.93–1.85), but with heterogeneity between studies ( $I^2 = 57.9\%$ ;  $P = 0.04$ ).

### Duration and severity of diabetes and NHL

There were limited and conflicting data on duration and severity of diabetes and risk of NHL, which were not uniformly reported, making formal statistical analyses impossible. Two studies reported a positive association between diabetes and risk of NHL with increasing duration of diabetes (5,18); however, in another study, the risk of NHL continuously decreased with increasing time of follow-up (20). In case-control studies, Fortuny et al. (24) showed an increased OR of NHL in individuals who had diabetes for

more than 12 years compared with patients without diabetes, whereas Scotti et al. (25) reported no increased risk of NHL with longer duration of diabetes.

### Diabetes therapy and NHL

There were limited data on therapy for diabetes and risk of NHL, which did not allow for formal statistical analyses. Cerhan et al. (18) reported an increased risk of NHL in women with diabetes who were on oral antidiabetic therapy (relative risk 3.43 [95% CI 1.67–7.07]), but the risk became nonstatistically significant in women who were on oral antidiabetic therapy and concurrent insulin (relative risk 2.13 [0.78–5.79]). Fortuny et al. (24) reported an increased risk in pa-

tients with diabetes not on pharmacological therapy (OR 1.73 [1.11–2.68]), whereas the risk in patients treated with oral antidiabetic therapy and/or insulin was not statistically significant.

**CONCLUSIONS**— Our analyses showed that patients with diabetes have an increased risk of developing NHL compared with those without diabetes. Although the size of the increased risk was moderate, based on the projected increase in the incidence and prevalence of diabetes (2), our findings suggest a concurrent increase in the incidence of NHL.

NHL is a malignant disorder characterized by an uncontrolled growth of monoclonal lymphocytes, observed most commonly within the lymph nodes. NHL is a heterogeneous condition and comprises several different morphologic subtypes with distinct clinical behaviors and outcomes. The increase in NHL incidence over the last few decades is part of a longer trend documented since 1950 (1), and improved medical technology and changing systems of diagnostic classification can explain only part of it. Many medical conditions have been associated with increased risk of developing lymphoma, including congenital or acquired immunodeficiency states (HIV [30] and post transplant [31]), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, and Sjögren's Syndrome) (32,33), and different infectious agents (34) such as human T-cell lymphotropic virus type I, hepatitis C virus, *Helicobacter pylori*, *Chlamydia psittaci*, and *Borrelia sp.* What most of these conditions have in common is immune dysregulation (9). Indeed, the risk of developing NHL among patients in immunosuppressed states is several times greater than that of the general population (30,31). However, these immunocompromized states are relatively rare and cannot fully explain the long-term and large increase in NHL incidence. Our findings of an increased risk of NHL associated with diabetes may provide an explanation for the increased incidence of NHL not associated with rare immunosuppressed states. It is increasingly recognized that not only type 1 but also type 2 diabetes is characterized by immune dysfunction related to impaired neutrophil activity and changes in cellular and humoral immunity (35), which may, at least in part, account for the increased risk of NHL that we found in this study.

The majority of patients with type 2 diabetes are obese; therefore, the observed association between diabetes and NHL may be mediated by metabolic derangements associated with the obese state. It is increasingly recognized that excess body fat is associated with increased risk of common and less common malignancies (36). A recent meta-analysis also found an increased risk of NHL with excess body weight (37). The increase in NHL associated with excess body weight might be mediated by changes in circulating adipocytokines, such as adiponectin and leptin. These cytokines are thought to be involved in immunity and inflammation and can alter the balance between cell proliferation and death. Studies in vitro and in animal models indicate that adiponectin has anti-inflammatory properties and reduces cell proliferation, whereas leptin has proinflammatory properties and promotes the growth of certain cancer cells. In vitro studies have shown that leptin stimulates the proliferation of normal hematopoietic cells as well as circulating monocytes producing proinflammatory cytokines, such as tumor-necrosis factor and interleukin.

Treatment of diabetes, rather than the disease pathophysiology itself, may also be of etiologic importance. For example, adverse reactions to sulfonylureas include agranulocytosis, aplastic anemia, bone marrow aplasia, and hemolytic anemia, but these clinically apparent conditions are relatively rare, and tumor-inducing effects have not been reported. Furthermore, various types of insulin are also known to be immunogenic. Based on the reviewed studies, we were unable to reach any conclusions on a potential link between therapy and NHL risk. Also, any potential treatment effects may be confounded by duration of the disease.

The increased risk of NHL in patients with diabetes was observed in both prospective and case-control studies but was statistically significant only among prospective studies. The disparity may be attributed to a variety of study-specific factors, including the relatively small number of cases, the failure to take confounding factors into account, and recall bias that may have resulted in misclassifications of type and duration of diabetes.

Our subgroup analysis showed that the risk of NHL in women with diabetes is 38% higher compared with women without diabetes, but the risk in men was not increased. These findings are in contrast with the reported increased ratio of

lymphoproliferative disorders in men compared with women. Although our findings may represent a true association between diabetes and NHL that occurs only in women, our results may be limited by the small number of studies that reported data by sex.

Based on the age of diagnosis as a proxy for type of diabetes, the risk of NHL appeared to be higher in patients with type 2 diabetes (older than 30 years) than in those with type 1 diabetes (younger than 30 years). However, the results for neither subgroup analysis were statistically significant.

Our study has certain limitations based on the quality of the published studies. First, although some studies confirmed the diagnosis of diabetes, most used self-report of diabetes as the predictor. A positive self-report for diabetes is generally quite accurate in epidemiologic studies (38). Second, the type of diabetes was not confirmed. Most studies used age as a proxy criterion for the two most common types of diabetes (type 1 and type 2). Severity, duration, and treatment of diabetes were also not reported in most studies. Third, our observed association may be due to confounding, as most prospective studies did not control for potential confounders, including HIV status of participants. Finally, our systematic review was restricted to articles in the English language, which may have influenced the results. However, there is also evidence that some countries publish unusually high proportions of positive results (39), which would introduce publication bias.

In conclusion, our analysis shows that diabetes is associated with moderately increased risk of NHL, which is consistent with other reported associations between diabetes and malignancies. Although the relative risk is moderate, given the rapidly increasing incidence and prevalence of diabetes, the number of incident cases of NHL attributed to diabetes can potentially be very high. Future studies should focus on elucidating potential pathophysiologic links between diabetes and NHL.

**Acknowledgments**—A.G.P. is supported by National Institutes of Health grants R01-DK76092, R0179003, and R2178867.

Portions of the manuscript were presented at the 68th Scientific Sessions of the American Diabetes Association, San Francisco, California, 6–10 June 2008.

## References

1. Chiu BC, Weisenburger DD: An update of the epidemiology of non-Hodgkin's lymphoma. *Clin Lymphoma* 4:161–168, 2003
2. Winer N, Sowers JR: Epidemiology of diabetes. *J Clin Pharmacol* 44:397–405, 2004
3. El-Serag HB, Hampel H, Javadi F: The association between diabetes and hepatocellular carcinoma: a systematic review of epidemiologic evidence. *Clin Gastroenterol Hepatol* 4:369–380, 2006
4. Everhart J, Wright D: Diabetes mellitus as a risk factor for pancreatic cancer. A meta-analysis. *Jama* 273:1605–1609, 1995
5. Hjalgrim H, Frisch M, Ekblom A, Kyvik KO, Melbye M, Green A: Cancer and diabetes: a follow-up study of two population-based cohorts of diabetic patients. *J Intern Med* 241:471–475, 1997
6. Khan M, Mori M, Fujino Y, Shibata A, Sakauchi F, Washio M, Tamakoshi A: Site-specific cancer risk due to diabetes mellitus history: evidence from the Japan Collaborative Cohort (JACC) Study. *Asian Pac J Cancer Prev* 7:253–259, 2006
7. Rousseau MC, Parent ME, Pollak MN, Siemiatycki J: Diabetes mellitus and cancer risk in a population-based case-control study among men from Montreal, Canada. *Int J Cancer* 118:2105–2109, 2006
8. Pickup JC: Inflammation and activated innate immunity in the pathogenesis of type 2 diabetes. *Diabetes Care* 27:813–823, 2004
9. Grulich AE, Vajdic CM, Cozen W: Altered immunity as a risk factor for non-Hodgkin lymphoma. *Cancer Epidemiol Biomarkers Prev* 16:405–408, 2007
10. Frasca F, Pandini G, Sciacca L, Pezzino V, Squatrito S, Belfiore A, Vigneri R: The role of insulin receptors and IGF-I receptors in cancer and other diseases. *Arch Physiol Biochem* 114:23–37, 2008
11. Lisker SA, Brody JL, Beizer LH: Abnormal carbohydrate metabolism in patients with malignant blood dyscrasias. *Am J Med Sci* 252:282–288, 1966
12. Harris NL, Jaffe ES, Diebold J, Flandrin G, Muller-Hermelink HK, Vardiman J, Lister TA, Bloomfield CD: World Health Organization classification of neoplastic diseases of the hematopoietic and lymphoid tissues: report of the Clinical Advisory Committee meeting-Airlie House, Virginia, November 1997. *J Clin Oncol* 17:3835–3849, 1999
13. DerSimonian R, Laird N: Meta-analysis in clinical trials. *Control Clin Trials* 7:177–188, 1986
14. Odds ratio versus relative risk [Doctoring Curriculum. Medical University of South Carolina], [article online]. Available from <http://www.musc.edu/dc/icrebm/oddsratio.html>. Accessed 10 May 2008
15. Lau J, Ioannidis JP, Schmid CH: Quantitative synthesis in systematic reviews. *Ann*

- Intern Med* 127:820–826, 1997
16. Higgins JP, Thompson SG, Deeks JJ, Altman DG: Measuring inconsistency in meta-analyses. *Bmj* 327:557–560, 2003
  17. Bax L, Yu LM, Ikeda N, Tsuruta H, Moons KG: Development and validation of MIX: comprehensive free software for meta-analysis of causal research data. *BMC Med Res Methodol* 6:50, 2006
  18. Cerhan JR, Wallace RB, Folsom AR, Potter JD, Sellers TA, Zheng W, Lutz CT: Medical history risk factors for non-Hodgkin's lymphoma in older women. *J Natl Cancer Inst* 89:314–318, 1997
  19. Swerdlow AJ, Laing SP, Qiao Z, Slater SD, Burden AC, Botha JL, Waugh NR, Morris AD, Gatling W, Gale EA, Patterson CC, Keen H: Cancer incidence and mortality in patients with insulin-treated diabetes: a UK cohort study. *Br J Cancer* 92:2070–2075, 2005
  20. Weiderpass E, Gridley G, Ekblom A, Nyren O, Hjalgrim H, Adami HO: Medical history risk factors for non-Hodgkin's lymphoma in older women. *J Natl Cancer Inst* 89:816–817, 1997
  21. Cerhan JR, Bernstein L, Severson RK, Davis S, Colt JS, Blair A, Hartge P: Anthropometrics, physical activity, related medical conditions, and the risk of non-hodgkin lymphoma. *Cancer Causes Control* 16:1203–1214, 2005
  22. Zahm SH, Blair A, Cantor KP, Fraumeni JF Jr.: Non-insulin dependent diabetes mellitus and non-Hodgkin's lymphoma. Other American studies fail to confirm an association. *BMJ* 310:1009–1010, 1995
  23. Cartwright RA, McKinney PA, O'Brien C, Richards ID, Roberts B, Lauder I, Darwin CM, Bernard SM, Bird CC: Non-Hodgkin's lymphoma: case control epidemiological study in Yorkshire. *Leuk Res* 12:81–88, 1988
  24. Fortuny J, Benavente Y, Bosch R, Garcia-Villanueva M, de Sevilla AF, de Sanjose S: Type 2 diabetes mellitus, its treatment and risk for lymphoma. *Eur J Cancer* 41: 1782–1787, 2005
  25. Scotti L, Tavani A, Bosetti C, Maso LD, Talamini R, Montella M, Franceschi S, La Vecchia C: Diabetes and risk of non-Hodgkin lymphoma: a case-control study. *Tumori* 93:1–3, 2007
  26. Vineis P, Crosignani P, Sacerdote C, Fontana A, Masala G, Miligi L, Nanni O, Ramazzotti V, Rodella S, Stagnaro E, Tumino R, Vigano C, Vindigni C, Costantini AS: Haematopoietic cancer and medical history: a multicentre case control study. *J Epidemiol Community Health* 54: 431–436, 2000
  27. Lin SY, Hsieh MS, Chen LS, Chiu YH, Yen AM, Chen TH: Diabetes mellitus associated with the occurrence and prognosis of non-Hodgkin's lymphoma. *Eur J Cancer Prev* 16:471–478, 2007
  28. Bernstein L, Ross RK: Prior medication use and health history as risk factors for non-Hodgkin's lymphoma: preliminary results from a case-control study in Los Angeles County. *Cancer Res* 52:S5510–S5515, 1992
  29. Smedby KE, Hjalgrim H, Askling J, Chang ET, Gregersen H, Porwit-MacDonald A, Sundstrom C, Akerman M, Melbye M, Glimelius B, Adami HO: Autoimmune and chronic inflammatory disorders and risk of non-Hodgkin lymphoma by subtype. *J Natl Cancer Inst* 98:51–60, 2006
  30. Cheung MC, Pantanowitz L, Dezube BJ: AIDS-related malignancies: emerging challenges in the era of highly active anti-retroviral therapy. *Oncologist* 10:412–426, 2005
  31. Andreone P, Gramenzi A, Lorenzini S, Biselli M, Cursaro C, Pileri S, Bernardi M: Posttransplantation lymphoproliferative disorders. *Arch Intern Med* 163:1997–2004, 2003
  32. Ekstrom Smedby K, Vajdic CM, Falster M, Engels EA, Martinez-Maza O, Turner J, Hjalgrim H, Vineis P, Seniori Costantini A, Bracci PM, Holly EA, Willett E, Spinelli JJ, La Vecchia C, Zheng T, Becker N, De Sanjose S, Chiu BC, Dal Maso L, Cocco P, Maynadie M, Foretova L, Staines A, Brennan P, Davis S, Severson R, Cerhan JR, Breen EC, Birmann B, Grulich AE, Cozen W: Autoimmune disorders and risk of non-Hodgkin lymphoma subtypes: a pooled analysis within the InterLymph Consortium. *Blood* 111:4029–4038, 2008
  33. Zintzaras E, Voulgarelis M, Moutsopoulos HM: The risk of lymphoma development in autoimmune diseases: a meta-analysis. *Arch Intern Med* 165:2337–2344, 2005
  34. Engels EA: Infectious agents as causes of non-Hodgkin lymphoma. *Cancer Epidemiol Biomarkers Prev* 16:401–404, 2007
  35. Moutschen MP, Scheen AJ, Lefebvre PJ: Impaired immune responses in diabetes mellitus: analysis of the factors and mechanisms involved. Relevance to the increased susceptibility of diabetic patients to specific infections. *Diabetes Metab* 18: 187–201, 1992
  36. Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M: Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet* 371:569–578, 2008
  37. Larsson SC, Wolk A: Obesity and risk of non-Hodgkin's lymphoma: a meta-analysis. *Int J Cancer* 121:1564–1570, 2007
  38. Midthjell K, Holmen J, Bjorndal A, Lund-Larsen G: Is questionnaire information valid in the study of a chronic disease such as diabetes? The Nord-Trondelag diabetes study. *J Epidemiol Community Health* 46: 537–542, 1992
  39. Vickers A, Goyal N, Harland R, Rees R: Do certain countries produce only positive results? A systematic review of controlled trials. *Control Clin Trials* 19:159–166, 1998